

# REFINED ENUMERATIONS OF ALTERNATING SIGN MATRICES: MONOTONE $(d, m)$ -TRAPEZOIDS WITH PRESCRIBED TOP AND BOTTOM ROW

ILSE FISCHER

**ABSTRACT.** Monotone triangles are plane integer arrays of triangular shape with certain monotonicity conditions along rows and diagonals. Their significance is mainly due to the fact that they correspond to  $n \times n$  alternating sign matrices when prescribing  $(1, 2, \dots, n)$  as bottom row of the array. We define monotone  $(d, m)$ -trapezoids as monotone triangles with  $m$  rows where the  $d - 1$  top rows are removed. (These objects are also equivalent to certain partial alternating sign matrices.) It is known that the number of monotone triangles with bottom row  $(k_1, \dots, k_n)$  is given by a polynomial  $\alpha(n; k_1, \dots, k_n)$  in the  $k_i$ 's. The main purpose of this paper is to show that the number of monotone  $(d, m)$ -trapezoids with prescribed top and bottom row appears as a coefficient in the expansion of a specialization of  $\alpha(n; k_1, \dots, k_n)$  with respect to a certain polynomial basis. This settles a generalization of a recent conjecture of Romik and the author. Among other things, the result is used to express the number of monotone triangles with bottom row  $(1, 2, \dots, i - 1, i + 1, \dots, j - 1, j + 1, \dots, n)$  (which is, by the standard bijection, also the number of  $n \times n$  alternating sign matrices with given top two rows) in terms of the number of  $n \times n$  alternating sign matrices with prescribed top and bottom row, and, by a formula of Stroganov for the latter numbers, to provide an explicit formula for the first numbers. (A formula of this type was first derived by Karklinsky and Romik using the relation of alternating sign matrices to the six-vertex model.)

## 1. INTRODUCTION

A *monotone triangle* is a triangular array of integers of the following form

$$\begin{array}{cccccccc}
& & & & & & & a_{n,n} \\
& & & & & & a_{n-1,n-1} & a_{n-1,n} \\
& & & \dots & & \dots & & \dots \\
& & a_{3,3} & & \dots & & \dots & a_{3,n} \\
& a_{2,2} & & a_{2,3} & & \dots & & \dots & a_{2,n} \\
a_{1,1} & & a_{1,2} & & a_{1,3} & & \dots & \dots & a_{1,n}
\end{array},$$

which is monotone increasing in northeast and in southeast direction and strictly increasing along rows, that is  $a_{i,j} \leq a_{i+1,j+1}$  for  $1 \leq i \leq j < n$ ,  $a_{i,j} \leq a_{i-1,j}$  for  $1 < i \leq j \leq n$  and  $a_{i,j} < a_{i,j+1}$  for  $1 \leq i \leq j \leq n - 1$ . Monotone triangles with bottom row  $(1, 2, \dots, n)$  are said to be *complete* and correspond to  $n \times n$  alternating sign matrices. The latter are defined as square matrices with entries in  $\{0, 1, -1\}$  such that in each row and column the sum of entries is 1 and the non-zero entries alternate. Next we display a complete monotone triangle and its corresponding alternating sign matrix.

---

Supported by the Austrian Science Foundation (FWF), grant number S9607-N13, in the framework of the National Research Network ‘‘Analytic Combinatorics and Probabilistic Number Theory’’.

$$\begin{array}{cccccc}
& & & 4 & & \\
& & & 3 & 4 & \\
& & 2 & 3 & 5 & \\
& 1 & 2 & 4 & 6 & \\
1 & 2 & 3 & 4 & 5 & 6
\end{array}
\begin{pmatrix} 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & -1 & 1 & 0 \\ 1 & 0 & -1 & 1 & -1 & 1 \\ 0 & 0 & 1 & -1 & 1 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \end{pmatrix}$$

In general, for a given complete monotone triangle  $A = (a_{i,j})_{1 \leq i \leq j \leq n}$  with  $n$  rows, the corresponding  $n \times n$  alternating sign matrix  $M = (m_{i,j})_{1 \leq i,j \leq n}$  can be obtained as follows: we have  $m_{i,j} = 1$  if and only if

$$j \in \{a_{n+1-i,n+1-i}, a_{n+1-i,n+2-i}, \dots, a_{n+1-i,n}\} \setminus \{a_{n+2-i,n+2-i}, a_{n+2-i,n+3-i}, \dots, a_{n+2-i,n}\},$$

$m_{i,j} = -1$  if and only if

$$j \in \{a_{n+2-i,n+2-i}, a_{n+2-i,n+3-i}, \dots, a_{n+2-i,n}\} \setminus \{a_{n+1-i,n+1-i}, a_{n+1-i,n+2-i}, \dots, a_{n+1-i,n}\}$$

and  $m_{i,j} = 0$  else.

The story of alternating sign matrices [1] started back in the 1980s when Mills, Robbins and Rumsey [7] conjectured that the number of  $n \times n$  alternating sign matrices is given by the following beautiful product formula

$$\prod_{j=0}^{n-1} \frac{(3j+1)!}{(n+j)!}.$$

It was more than 10 years later when Zeilberger [9] finally succeeded in giving the first proof of this formula. Since then refined enumerations of alternating sign matrices and enumerations of symmetry classes of alternating sign matrices have been accomplished and fruitful relations to other areas such as statistical physics (six-vertex model) and algebra have been discovered and made the combinatorial analysis of these objects to a major topic in enumerative and algebraic combinatorics as well as in statistical physics.

The central objects of this article are the following generalizations of monotone triangles: for  $1 \leq d \leq m$ , a *monotone  $(d, m)$ -trapezoid* is a monotone triangle with  $m$  rows where the first  $d-1$  rows are removed. Thus,

$$\begin{array}{cccccc}
& & 2 & 3 & 5 & \\
& 1 & 2 & 4 & 6 & \\
1 & 2 & 3 & 5 & 6 &
\end{array}$$

is a monotone  $(3, 5)$ -trapezoid. It is easy to translate these objects into the language of alternating sign matrices: for  $t \leq n$ , let a  $(t, n)$ -partial alternating sign matrix be a  $t \times n$  matrix with entries in  $\{0, 1, -1\}$  such that the non-zero entries alternate in each row and column and the rowsums are equal to 1. Suppose  $1 \leq i_1 < i_2 < \dots < i_d \leq n$  and  $1 \leq s_1 < s_2 < \dots < s_c \leq n$ . Then  $(d, n-c)$ -trapezoids with top row  $i_1, \dots, i_d$  and whose bottom row consists of the numbers in  $\{1, 2, \dots, n\} \setminus \{s_1, s_2, \dots, s_c\}$ , arranged in increasing order, are equivalent to  $(n-c-d, n)$ -partial alternating sign matrices with the following properties: the  $j$ -th columnsum is 1 iff  $j \notin \{i_1, i_2, \dots, i_d, s_1, s_2, \dots, s_c\}$  and it is  $-1$  iff  $j \in \{i_1, \dots, i_d\} \cap \{s_1, \dots, s_c\}$ . Otherwise, the  $j$ -th columnsum is 0 and the first non-zero entry (if there is any) of this column is a  $-1$  iff  $j \in \{i_1, \dots, i_d\} \setminus \{s_1, \dots, s_c\}$ . The  $(2, 6)$ -partial alternating sign matrix that corresponds to

the monotone  $(3, 5)$ -trapezoid given above is

$$\begin{pmatrix} 1 & 0 & -1 & 1 & -1 & 1 \\ 0 & 0 & 1 & -1 & 1 & 0 \end{pmatrix}_1.$$

Let  $\alpha(n; k_1, \dots, k_n)$  denote the number of monotone triangles with bottom row  $(k_1, \dots, k_n)$ . In [4] it was shown that

$$\alpha(n; k_1, \dots, k_n) = \left( \prod_{1 \leq p < q \leq n} (\text{id} + E_{k_p} E_{k_q} - E_{k_q}) \right) \prod_{1 \leq i < j \leq n} \frac{k_j - k_i}{j - i}, \quad (1.1)$$

where  $E_x$  denotes the shift operator, defined as  $E_x p(x) = p(x + 1)$ . Moreover note that the product of operators in this formula is understood as the composition. In particular, this operator formula implies that  $\alpha(n; k_1, \dots, k_n)$  is a polynomial in  $(k_1, \dots, k_n)$  of degree no greater than  $n - 1$  in each  $k_i$ . For non-negative integers  $c, d$  with  $c + d \leq n$ , we consider the following expansion.

$$\begin{aligned} & \alpha(n; k_1, \dots, k_c, c + 1, c + 2, \dots, n - d, k_{n-d+1}, k_{n-d+2}, \dots, k_n) \\ &= \sum_{s_1=1}^n \sum_{s_2=1}^n \dots \sum_{s_c=1}^n \sum_{i_1=1}^n \sum_{i_2=1}^n \dots \sum_{i_d=1}^n A(n; s_1, s_2, \dots, s_c; i_1, \dots, i_d) \\ & \times (-1)^{s_1+s_2+\dots+s_c+c} \binom{k_1 - c - 1}{s_c - 1} \binom{k_2 - c - 1}{s_{c-1} - 1} \dots \binom{k_c - c - 1}{s_1 - 1} \\ & \times \binom{k_{n-d+1} - n + d - 2 + i_1}{i_1 - 1} \binom{k_{n-d+2} - n + d - 2 + i_2}{i_2 - 1} \dots \binom{k_n - n + d - 2 + i_d}{i_d - 1}. \end{aligned} \quad (1.2)$$

The main subject of this paper is the following generalization of Conjecture 7 from [5], which provides a combinatorial interpretation of the coefficients  $A(n; s_1, s_2, \dots, s_c; i_1, \dots, i_d)$ . (In fact, this theorem also implies Conjecture 5 from [5], see Section 6.)

**Theorem 1.** *For  $1 \leq s_1 < s_2 < \dots < s_c \leq n$  and  $1 \leq i_1 < i_2 < \dots < i_d \leq n$  the coefficient  $A(n; s_1, \dots, s_c; i_1, \dots, i_d)$  is the number of monotone  $(d, n - c)$ -trapezoids with  $(i_1, \dots, i_d)$  as top row and whose bottom row consists of the numbers in  $\{1, 2, \dots, n\} \setminus \{s_1, s_2, \dots, s_c\}$ , arranged in increasing order.*

Its proof is the content of Sections 2–4. In Section 5, we use the identity

$$\alpha(n; k_1, \dots, k_n) = (-1)^{n-1} \alpha(n; k_2, \dots, k_n, k_1 - n)$$

from [3, Lemma 5] to express the coefficient

$$A(n; s_1, \dots, s_c; i_1, \dots, i_d)$$

in terms of the coefficients

$$A(n; s_1, \dots, s_{c-t}; i_1, \dots, i_{d+t})$$

where  $t$  is a fixed non-negative integer no greater than  $c$  and  $s_{c-l+1} \leq i_{d+l} \leq n$  for  $1 \leq l \leq t$ . In the special case  $c = 2$ ,  $d = 0$  and  $t = 1$  this relates  $A(n; s_1, s_2; -) =: A_{n, s_1, s_2}$  to  $A(n; s_1; i_1) =:$

---

<sup>1</sup>As the monotone  $(3, 5)$ -trapezoid consists of rows 3–5 of our introductory example of a complete monotone triangle, the  $(2, 6)$ -partial alternating sign matrix is the fourth and fifth row of the corresponding alternating sign matrix.

$B_{n,s_1,i_1}$ . By the standard bijection between alternating sign matrices and complete monotone triangles,  $B_{n,i,j}$  is the number of  $n \times n$  alternating sign matrices where the unique 1 in the bottom row is in column  $i$  and the unique 1 in the top row is in column  $j$ . Stroganov [8] has derived a formula for this doubly refined enumeration of alternating sign matrices. On the other hand, the numbers  $A_{n,i,j}$  have been studied in [5] and our relation to Stroganov's doubly refined enumeration numbers finally enables us to provide the following formula for the numbers  $A_{n,i,j}$ :

$$A_{n,i,j} = \frac{1}{A_{n-1}} \sum_{i=1}^l \sum_{k=l-i+j}^{l-i+n} (-1)^{n+i+k+l} \binom{2n-2-j}{k-l+i-j} \times (A_{n-1,l-1}(A_{n,k} - A_{n,k-1}) + A_{n-1,k-1}(A_{n,l} - A_{n,l-1})), \quad (1.3)$$

where

$$A_{n,k} = \begin{cases} \binom{n+k-2}{n-1} \frac{(2n-k-1)!}{(n-k)!} \prod_{j=0}^{n-2} \frac{(3j+1)!}{(n+j)!} & 1 \leq k \leq n \\ 0 & \text{otherwise} \end{cases}$$

is the number of  $n \times n$  alternating sign matrices that have a 1 in the  $k$ -th column of the first row and  $A_n$  denotes the total number of  $n \times n$  alternating sign matrices. A formula similar to (1.3) was first derived by Karklinsky and Romik [6] by using the six-vertex model approach to alternating sign matrices. Another, complicated, formula for  $A_{n,i,j}$  was conjectured in [5, Conjecture 4].)

In fact, the numbers  $A_{n,i,j}$  also correspond to a certain doubly refined enumeration of  $n \times n$  alternating sign matrices with respect to the two bottom (or, equivalently, two top) rows if  $i < j$ : let  $k$  be a fixed integer with  $i \leq k \leq j$ . Then  $A_{n,i,j}$  is the number of  $n \times n$  alternating sign matrices such that  $e_i + e_j - e_k$  is the  $(n-1)$ -st row and  $e_k$  is the bottom row where  $e_p \in \mathbb{R}^n$  with  $(e_p)_q = \delta_{p,q}$ . Alternatively and as explained above, it is also the number of  $(n-2, n)$ -partial alternating sign matrices such that the last non-zero entries (if there exist any) in columns  $i$  and  $j$  are  $-1$ s.

## 2. HOW DOES ROTATING AND REFLECTING ALTERNATING SIGN MATRICES TRANSLATE INTO THE LANGUAGE OF COMPLETE MONOTONE TRIANGLES?

The set of  $n \times n$  alternating sign matrices is invariant under the rotations of  $90^\circ$  and under the reflection along any of the four symmetry axes of the square. Here, we investigate how these rotations and reflections translate into the language of complete monotone triangles.

Let  $A = (a_{i,j})_{1 \leq i \leq j \leq n}$  be a complete monotone triangle with  $n$  rows and  $M$  be the corresponding  $n \times n$  alternating sign matrix. We fix the following notation: the sequence  $(a_{l,l}, a_{l-1,l}, \dots, a_{1,l})$  is said to be the  $l$ -th SE-diagonal and the sequence  $(a_{1,l}, a_{2,l+1}, \dots, a_{n-l+1,n})$  is said to be the  $l$ -th NE-diagonal of the complete monotone triangle  $A$ .

It is not hard to convince oneself that the complete monotone triangle  $B = (b_{i,j})_{1 \leq i \leq j \leq n}$  which corresponds to the alternating sign matrix that we obtain by reflecting  $M$  along the antidiagonal is given by

$$b_{i,j} = \# \text{ of elements } x \text{ in the } j\text{-th SE-diagonal of } A \text{ with } x \geq i.$$

We set  $B = \text{AD}(A)$ . For the example in the introduction we obtain the following monotone triangle and alternating sign matrix.

$$\begin{array}{cccccc}
& & & 3 & & \\
& & & 2 & 4 & \\
& & 1 & 3 & 6 & \\
& 1 & 2 & 5 & 6 & \\
1 & 2 & 4 & 5 & 6 & \\
1 & 2 & 3 & 4 & 5 & 6
\end{array}
\begin{pmatrix} 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 1 & -1 & 1 & 0 & 0 \\ 1 & -1 & 1 & -1 & 0 & 1 \\ 0 & 1 & -1 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \end{pmatrix}$$

Similarly, the complete monotone triangle  $C = (c_{i,j})_{1 \leq i \leq j \leq n}$  which corresponds to the alternating sign matrix that we obtain by rotating  $M$  clockwise by  $90^\circ$  is given by

$$c_{i,j} = \# \text{ of elements } x \text{ in the } (n+1-j)\text{-th NE-diagonal of } A \text{ with } x \leq n+1-i.$$

We set  $C = \text{ROT}(A)$ . In the example this gives the following two objects.

$$\begin{array}{cccccc}
& & & 3 & & \\
& & & 3 & 4 & \\
& & 2 & 4 & 5 & \\
& 1 & 3 & 5 & 6 & \\
1 & 2 & 4 & 5 & 6 & \\
1 & 2 & 3 & 4 & 5 & 6
\end{array}
\begin{pmatrix} 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & -1 & 0 & 1 & 0 \\ 1 & -1 & 1 & -1 & 0 & 1 \\ 0 & 1 & -1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \end{pmatrix}$$

Finally, the complete monotone triangle  $D = (d_{i,j})_{1 \leq i \leq j \leq n}$  that corresponds to the alternating sign matrix that we obtain by reflecting  $M$  along the horizontal symmetry axis is uniquely determined by

$$\{a_{i,i}, a_{i,i+1}, \dots, a_{i,n}, d_{n+2-i,n+2-i}, d_{n+2-i,n+3-i}, \dots, d_{n+2-i,n}\} = \{1, 2, \dots, n\}$$

for all  $i \in \{1, 2, \dots, n\}$ . Here we set  $D = \text{H}(A)$ . In this case the running example changes to

$$\begin{array}{cccccc}
& & & 4 & & \\
& & & 3 & 5 & \\
& & 1 & 4 & 6 & \\
& 1 & 2 & 5 & 6 & \\
1 & 2 & 3 & 5 & 6 & \\
1 & 2 & 3 & 4 & 5 & 6
\end{array}
\begin{pmatrix} 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & -1 & 1 & 0 \\ 1 & 0 & -1 & 1 & -1 & 1 \\ 0 & 1 & 0 & -1 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \end{pmatrix}.$$

Since reflecting an alternating sign matrix along the antidiagonal is equivalent to first rotating it clockwise by  $90^\circ$  and then reflecting it along the horizontal symmetry axis we have

$$\text{AD} = \text{H} \circ \text{ROT}. \quad (2.1)$$

However, in the following we do not need the interpretations of the mappings  $\text{AD}$ ,  $\text{ROT}$  and  $\text{H}$  in terms of alternating sign matrices. For our purpose, it suffices to show that these three mappings are permutations of the set of complete monotone triangles with  $n$  rows (this is easy and left to the reader) and that they satisfy (2.1). The latter is equivalent to showing that for every complete monotone triangle  $A = (a_{i,j})_{1 \leq i \leq j \leq n}$  and  $i \in \{1, 2, \dots, n\}$  the union of

$$\{\# \text{ of elements } x \text{ in } j\text{-th SE-diagonal of } A \text{ with } x \geq i \mid i \leq j \leq n\}$$

and

$$\{\# \text{ of elements } x \text{ in the } j\text{-th NE-diagonal of } A \text{ with } x \leq i-1 \mid 1 \leq j \leq i-1\}$$

is equal to  $\{1, 2, \dots, n\}$ . This follows from the following fact which can be shown by induction with respect to  $n$ , see Figure 1: suppose  $(p_1, p_2, \dots, p_{i-1})$  is a strictly decreasing sequence of

positive integers and for  $1 \leq j \leq i-1$  delete the first  $p_j$  entries from the  $j$ -th NE-diagonal of a monotone triangle with  $n$  rows. For  $i \leq j \leq n$ , let  $q_j$  be the length of the  $j$ -th SE-diagonal in the remaining (partial) monotone triangle. Then

$$\{p_1, \dots, p_{i-1}, q_i, q_{i+1}, \dots, q_n\} = \{1, 2, \dots, n\}.$$





FIGURE 1.  $p_1 = 5, p_2 = 4, p_3 = 2, p_4 = 1, q_5 = 3, q_6 = 6$


### 3. FROM MONOTONE $(d, n-c)$ -TRAPEZOIDS WITH PRESCRIBED TOP AND BOTTOM ROW TO OTHER PARTIAL MONOTONE TRIANGLES

In this section we fix the two sequences  $(i_1, \dots, i_d)$  and  $(s_1, \dots, s_c)$  with  $1 \leq i_1 < i_2 < \dots < i_d \leq n$  and  $1 \leq s_1 < s_2 < \dots < s_c \leq n$ . We will use the mappings from the previous section to show that the number of monotone  $(d, n-c)$ -trapezoids with  $(i_1, \dots, i_d)$  as top row and whose bottom row consists of the numbers in  $\{1, 2, \dots, n\} \setminus \{s_1, s_2, \dots, s_c\}$ , arranged in increasing order, is equal to the number of (partial) monotone triangles with  $n$  rows, entries in  $\{c+1, c+2, \dots, n-d\}$  where for  $1 \leq l \leq c$  the first  $s_{c+1-l}$  entries of the  $l$ -th NE-diagonal are missing and for  $n-d+1 \leq l \leq n$  the last  $i_{l-n+d}$  entries of the  $l$ -th SE-diagonal are missing.

Indeed, fix such a monotone  $(d, n-c)$ -trapezoid and add entries arbitrarily to make it up to a complete monotone triangle  $A$  with  $n$  rows. As an example, we take the  $(3, 5)$ -trapezoid on the left and the completion on the right, i.e.  $c = 1$  and  $n = 6$ .



Consider  $B = AD^{-1}(A)$ . In our example  $B$  is the following complete monotone triangle.



(The entries  $c+1, c+2, \dots, n-d$  are displayed in boldface as they correspond to the original monotone  $(3, 5)$ -trapezoid.) By the definition of  $AD$ , the elements greater than  $n-d$  in  $B$  are just the last  $i_{l-n+d}$  elements in the  $l$ -th SE-diagonals where  $n-d+1 \leq l \leq n$  and the distribution

of these entries is exactly determined by the completion of the monotone  $(d, n - c)$ -trapezoid towards the top. On the other hand, by (2.1),  $B = \text{ROT}^{-1} \circ H^{-1}(A)$ . Thus, the  $c$ -th row of  $H^{-1}(A)$  (counted from the top) is  $(s_1, \dots, s_c)$  and the first  $c - 1$  rows of this complete monotone triangle correspond to the completion of the original monotone  $(d, n - c)$ -trapezoid towards the bottom. Consequently (and by the definition of  $\text{ROT}$ ), the elements no greater than  $c$  of  $B$  are just the first  $s_{c+1-l}$  elements in the  $l$ -th NE-diagonals for  $1 \leq l \leq c$  and the distribution of these entries is determined by the completion of the monotone  $(d, n - c)$ -trapezoid towards the bottom.

If we delete in our example the first  $s_{c+1-l}$  entries in the  $l$ -th NE-diagonal for  $1 \leq l \leq c$  and the last  $i_{l-n+d}$  elements in the  $l$ -th SE-diagonal for  $n - d + 1 \leq l \leq n$  (i.e. exactly the part of the complete monotone triangles that corresponds to the completion of the original monotone  $(d, n - c)$ -trapezoid towards top and bottom), we clearly obtain the following partial monotone triangle.

$$\begin{array}{ccc} & & 3 \\ & 2 & 3 \\ & & 2 \\ & 2 & 3 \\ 2 & & 3 \end{array}$$

For technical reasons (which become apparent later) we add the entry  $c + 1$  at the beginning of the truncated  $l$ -th NE-diagonal for  $1 \leq l \leq c$  and add the entry  $n - d$  at the end of the truncated  $l$ -th SE-diagonal for  $n - d + 1 \leq l \leq n$ . In our example, we obtain

$$\begin{array}{ccc} & & 3 \\ & & 3 \\ & 2 & 3 \\ 2 & 2 & 3 \\ & 2 & 3 \\ 2 & 3 & 3 \end{array} \quad .$$

This will certainly not destroy the monotonicity along diagonals, but may disturb the strict increase along rows. All in all we see that the number of monotone  $(d, n - c)$ -trapezoids with fixed top and bottom row as given above is the number of partial monotone triangles with  $n$  rows and entries in  $\{c + 1, c + 2, \dots, n - d\}$ , where for  $1 \leq l \leq c$  the first  $s_{c+1-l} - 1$  entries of the  $l$ -th NE-diagonal are missing and the first entry of the truncated diagonal is equal to  $c + 1$  and does not necessarily have to be strictly smaller than its neighbour to the right and for  $n - d + 1 \leq l \leq n$  the last  $i_{l-n+d} - 1$  entries in the  $l$ -th SE-diagonal are missing and the last entry of the truncated diagonal is equal to  $n - d$  and does not necessarily have to be strictly greater than its neighbour to the left.

#### 4. COMPLETION OF THE PROOF OF THEOREM 1

Suppose  $A(l_1, \dots, l_{n-1})$  is a function in  $(l_1, \dots, l_{n-1})$  and  $(k_1, \dots, k_n) \in \mathbb{Z}^n$ . We define a summation operator

$$\sum_{(l_1, \dots, l_{n-1})}^{(k_1, \dots, k_n)} A(l_1, \dots, l_{n-1})$$

by induction with respect to  $n$ . If  $n = 0$  then the application of the operator gives zero, for  $n = 1$  we set  $\sum_{(-)}^{k_1} A = A$ . If  $n \geq 2$  then we define

$$\sum_{(l_1, \dots, l_{n-1})}^{(k_1, \dots, k_n)} A(l_1, \dots, l_{n-1}) = \sum_{(l_1, \dots, l_{n-2})}^{(k_1, \dots, k_{n-1})} \sum_{l_{n-1}=k_{n-1}}^{k_n} A(l_1, \dots, l_{n-1}) - \sum_{(l_1, \dots, l_{n-3})}^{(k_1, \dots, k_{n-2})} A(l_1, \dots, l_{n-3}, k_{n-1}, k_{n-1}).$$

With this, we obviously have

$$\alpha(n; k_1, \dots, k_n) = \sum_{(l_1, \dots, l_{n-1})}^{(k_1, \dots, k_n)} \alpha(n-1; l_1, \dots, l_{n-1})$$

for  $n \geq 2$ . In fact, if we extend the definition of monotone triangles to triangular integer arrays  $(a_{i,j})_{1 \leq i \leq j \leq n}$  with weak increase along NE-diagonals and SE-diagonals and strict increase along rows with the possible exception of the bottom row which may only be weakly increasing then  $\alpha(n; k_1, \dots, k_n)$  is also the number of these (extended) monotone triangles with bottom row  $(k_1, \dots, k_n)$ . Let  $\delta_x = \text{id} - E_x^{-1}$  and set

$$B(k_1, \dots, k_n) = \sum_{(l_1, \dots, l_{n-1})}^{(k_1, \dots, k_n)} A(l_1, \dots, l_{n-1}).$$

The recursion shows that

$$\delta_{k_n} B(k_1, \dots, k_n) = \sum_{(l_1, \dots, l_{n-2})}^{(k_1, \dots, k_{n-1})} A(l_1, \dots, l_{n-2}, k_n)$$

and, more general for  $d \geq 1$ ,

$$\delta_{k_{n-d+1}} \delta_{k_{n-d+2}} \dots \delta_{k_n} B(k_1, \dots, k_n) = \sum_{(l_1, \dots, l_{n-d-1})}^{(k_1, \dots, k_{n-d})} A(l_1, \dots, l_{n-d-1}, k_{n-d+1}, \dots, k_n).$$

Similarly, with  $\Delta_x = E_x - \text{id}$ , it is not hard to see that

$$(-1)^c \Delta_{k_1} \Delta_{k_2} \dots \Delta_{k_c} B(k_1, \dots, k_n) = \sum_{(l_{c+1}, \dots, l_{n-1})}^{(k_{c+1}, \dots, k_n)} A(k_1, \dots, k_c, l_{c+1}, \dots, l_{n-1}).$$

If we combine these two facts we see that

$$\begin{aligned} \Delta_{k_1} \Delta_{k_2} \dots \Delta_{k_c} \delta_{k_{n-d+1}} \delta_{k_{n-d+2}} \dots \delta_{k_n} (-1)^c B(k_1, \dots, k_n) \\ = \sum_{(l_{c+1}, \dots, l_{n-d-1})}^{(k_{c+1}, \dots, k_{n-d})} A(k_1, \dots, k_c, l_{c+1}, \dots, l_{n-d-1}, k_{n-d+1}, \dots, k_n). \end{aligned} \quad (4.1)$$






FIGURE 2. In this example we have  $n = 12$ ,  $c = d = 4$ ,  $s_1 = 2, s_2 = 4, s_3 = 7, s_4 = 10$  and  $i_1 = 3, i_2 = 6, i_3 = 8, i_4 = 9$ .

For  $1 \leq i_1 \leq i_2 \leq \dots \leq i_d \leq n$  and  $1 \leq s_1 \leq s_2 \leq \dots \leq s_c \leq n$ , let

$$\gamma(n; k_1, \dots, k_n; s_1, \dots, s_c; i_1, \dots, i_d)$$

(see Figure 4) denote the number of partial monotone triangles with  $n$  rows that have the following properties.

- For  $1 \leq l \leq c$ , the first  $s_{c+1-l} - 1$  entries of the  $l$ -th NE-diagonal are missing.
- The first entry of the truncated  $l$ -th NE-diagonal is equal to  $k_l$  and does neither have to be strictly smaller than its neighbour to the right nor weakly smaller than its SE-neighbour.
- For  $n - d + 1 \leq l \leq n$ , the last  $i_{l-n+d} - 1$  entries in the  $l$ -th SE-diagonal are missing.
- The last entry of the truncated  $l$ -th SE-diagonal is equal to  $k_l$  and does neither have to be strictly greater than its neighbour to the left nor weakly greater than its SW-neighbour.
- The bottom row (without those entries among  $k_1, k_2, \dots, k_c, k_{n-d+1}, k_{n-d+2}, \dots, k_n$  that are situated in the bottom row) is equal to  $k_{c+1}, k_{c+2}, \dots, k_{n-d}$  and we do not demand a strict increase for those entries.

We want to show that

$$\begin{aligned} & \gamma(n; k_1, \dots, k_n; s_1, \dots, s_c; i_1, \dots, i_d) \\ &= \Delta_{k_1}^{s_c-1} \Delta_{k_2}^{s_{c-1}-1} \dots \Delta_{k_c}^{s_1-1} \delta_{k_{n-d+1}}^{i_1-1} \delta_{k_{n-d+2}}^{i_2-1} \dots \delta_{k_n}^{i_d-1} (-1)^{s_1+\dots+s_c-c} \alpha(n; k_1, \dots, k_n) \end{aligned} \quad (4.2)$$

by induction with respect to  $n$ . The case  $n = 1$  is easy to check. Otherwise, let  $c' \geq 0$  be minimal such that  $s_{c'+1} > 1$  and  $d' \geq 0$  be minimal such that  $i_{d'+1} > 1$ . Then we have the

following recursion for  $\gamma$  with respect to  $n$ .

$$\gamma(n; k_1, \dots, k_n; s_1, \dots, s_c; i_1, \dots, i_d) = \sum_{(l_{c-c'+1}, \dots, l_{n-d+d'-1})}^{(k_{c-c'+1}, \dots, k_{n-d+d'})} \gamma(n-1; k_1, \dots, k_{c-c'}, l_{c-c'+1}, \dots, l_{n-d+d'-1}, k_{n-d+d'+1}, \dots, k_n; s_{c'+1}-1, s_{c'+2}-1, \dots, s_c-1; i_{d'+1}-1, i_{d'+2}-1, \dots, i_d-1)$$

By the induction hypothesis, this is equal to

$$\sum_{(l_{c-c'+1}, \dots, l_{n-d+d'-1})}^{(k_{c-c'+1}, \dots, k_{n-d+d'})} \Delta_{k_1}^{s_c-2} \Delta_{k_2}^{s_{c-1}-2} \dots \Delta_{k_{c-c'}}^{s_{c'+1}-2} \delta_{k_{n-d+d'+1}}^{i_{d'+1}-2} \dots \delta_{k_n}^{i_d-2} (-1)^{(s_{c'+1}-1)+\dots+(s_c-1)-c+c'} \alpha(n-1; k_1, \dots, k_{c-c'}, l_{c-c'+1}, \dots, l_{n-d+d'-1}, k_{n-d+d'+1}, \dots, k_n).$$

Now, (4.1) implies that this is furthermore equal to

$$\Delta_{k_1}^{s_c-2} \Delta_{k_2}^{s_{c-1}-2} \dots \Delta_{k_{c-c'}}^{s_{c'+1}-2} \delta_{k_{n-d+d'+1}}^{i_{d'+1}-2} \dots \delta_{k_n}^{i_d-2} (-1)^{s_{c'+1}+\dots+s_c} \Delta_{k_1} \Delta_{k_2} \dots \Delta_{k_{c-c'}} \delta_{k_{n-d+d'+1}} \delta_{k_{n-d+d'+2}} \dots \delta_{k_n} (-1)^{c-c'} \alpha(n; k_1, \dots, k_n).$$

This concludes the proof of (4.2).

By the observations from the previous section and (4.2), the number of  $(d, n-c)$ -trapezoids with  $(i_1, \dots, i_d)$  as top row and whose bottom row consists of the numbers in  $\{1, 2, \dots, n\} \setminus \{s_1, s_2, \dots, s_c\}$ , arranged in increasing order, is equal to

$$\Delta_{k_1}^{s_c-1} \Delta_{k_2}^{s_{c-1}-1} \dots \Delta_{k_c}^{s_1-1} \delta_{k_{n-d+1}}^{i_1-1} \delta_{k_{n-d+2}}^{i_2-1} \dots \delta_{k_n}^{i_d-1} \times (-1)^{s_1+\dots+s_c-c} \alpha(n; k_1, \dots, k_n) \Big|_{(k_1, \dots, k_n) = ((c+1)^c, c+1, c+2, \dots, n-d, (n-d)^d)}.$$

Finally, by the expansion (1.2), this number is equal to

$$\sum_{t_1=1}^n \dots \sum_{t_c=1}^n \sum_{j_1=1}^n \dots \sum_{j_d=1}^n A(n; t_1, \dots, t_c; j_1, \dots, j_d) (-1)^{t_1+\dots+t_c+s_1+\dots+s_c} \prod_{l=1}^c \Delta_{k_l}^{s_{c+1-l}-1} \binom{k_l - c - 1}{t_{c+1-l} - 1} \times \prod_{l=n-d+1}^n \delta_{k_l}^{i_{l-n+d}-1} \binom{k_l - n + d - 2 + j_{l-n+d}}{j_{l-n+d} - 1} \Big|_{(k_1, \dots, k_c, k_{n-d+1}, \dots, k_n) = ((c+1)^c, (n-d)^d)}.$$

Since  $\Delta_x \binom{x}{n} = \binom{x}{n-1}$  and  $\delta_x \binom{x}{n} = \binom{x-1}{n-1}$ , this is equal to

$$\begin{aligned}
 & \sum_{t_1=1}^n \cdots \sum_{t_c=1}^n \sum_{j_1=1}^n \cdots \sum_{j_d=1}^n A(n; t_1, \dots, t_c; j_1, \dots, j_d) (-1)^{t_1 + \cdots + t_c + s_1 + \cdots + s_c} \prod_{l=1}^c \binom{k_l - c - 1}{t_{c+1-l} - s_{c+1-l}} \\
 & \quad \times \prod_{l=n-d+1}^n \binom{k_l - n + d - 1 + j_{l-n+d} - i_{l-n+d}}{j_{l-n+d} - i_{l-n+d}} \Bigg|_{(k_1, \dots, k_c, k_{n-d+1}, \dots, k_n) = ((c+1)^c, (n-d)^d)} \\
 & = \sum_{t_1=1}^n \cdots \sum_{t_c=1}^n \sum_{j_1=1}^n \cdots \sum_{j_d=1}^n A(n; t_1, \dots, t_c; j_1, \dots, j_d) (-1)^{t_1 + \cdots + t_c + s_1 + \cdots + s_c} \\
 & \quad \times \prod_{l=1}^c \binom{0}{t_{c+1-l} - s_{c+1-l}} \prod_{l=n-d+1}^n \binom{j_{l-n+d} - i_{l-n+d} - 1}{j_{l-n+d} - i_{l-n+d}}
 \end{aligned}$$

As  $\binom{0}{k} = \delta_{0,k} = \binom{k-1}{k}$  for all integers  $k$ , this simplifies to  $A(n; s_1, \dots, s_c; i_1, \dots, i_d)$  and concludes the proof of Theorem 1.

## 5. A FORMULA FOR $A_{n,i,j}$

In order to prove (1.3) we make use of the identity

$$\alpha(n; k_1, \dots, k_n) = (-1)^{n-1} \alpha(n; k_2, \dots, k_n, k_1 - n),$$

which can be found in [3, Lemma 5]. It implies

$$\alpha(n; k_1, \dots, k_n) = (-1)^{tn-t} \alpha(n; k_{t+1}, \dots, k_n, k_1 - n, k_2 - n, \dots, k_t - n)$$

for  $t \geq 0$ . Moreover, for every integer  $z$ ,

$$\alpha(n; k_1, \dots, k_n) = \alpha(n; k_1 + z, \dots, k_n + z).$$

Therefore,

$$\begin{aligned}
& \sum_{s_1=1}^n \cdots \sum_{s_c=1}^n \sum_{i_1=1}^n \cdots \sum_{i_d=1}^n A(n; s_1, \dots, s_c; i_1, \dots, i_d) (-1)^{s_1+\dots+s_c+c} \\
& \quad \times \prod_{l=1}^c \binom{k_l - c - 1}{s_{c+1-l} - 1} \prod_{l=n-d+1}^n \binom{k_l - n + d - 2 + i_{l-n+d}}{i_{l-n+d} - 1} \\
& \quad = \alpha(n; k_1, \dots, k_c, c+1, c+2, \dots, n-d, k_{n-d+1}, \dots, k_n) \\
& \quad = (-1)^{tn-t} \alpha(n; k_{t+1}, \dots, k_c, c+1, c+2, \dots, n-d, k_{n-d+1}, \dots, k_n, k_1-n, k_2-n, \dots, k_t-n) \\
& \quad \quad = (-1)^{tn-t} \\
& \quad \times \alpha(n; k_{t+1}-t, \dots, k_c-t, c+1-t, c+2-t, \dots, n-d-t, k_{n-d+1}-t, \dots, k_n-t, k_1-n-t, \dots, k_t-n-t) \\
& \quad = \sum_{s_1=1}^n \cdots \sum_{s_{c-t}=1}^n \sum_{i_1=1}^n \cdots \sum_{i_{d+t}=1}^n A(n; s_1, \dots, s_{c-t}; i_1, \dots, i_{d+t}) (-1)^{s_1+\dots+s_{c-t}+c+tn} \\
& \quad \quad \times \binom{k_{t+1}-c-1}{s_{c-t}-1} \binom{k_{t+2}-c-1}{s_{c-t-1}-1} \cdots \binom{k_c-c-1}{s_1-1} \\
& \quad \quad \times \binom{k_{n-d+1}-n+d-2+i_1}{i_1-1} \cdots \binom{k_n-n+d-2+i_d}{i_d-1} \\
& \quad \quad \left( \binom{k_1-2n+d-2+i_{d+1}}{i_{d+1}-1} \binom{k_2-2n+d-2+i_{d+2}}{i_{d+2}-1} \cdots \binom{k_t-2n+d-2+i_{d+t}}{i_{d+t}-1} \right).
\end{aligned}$$

This implies that

$$\begin{aligned}
& \sum_{s_{c-t+1}=1}^n \sum_{s_{c-t+2}=1}^n \cdots \sum_{s_c=1}^n A(n; s_1, \dots, s_c; i_1, \dots, i_d) (-1)^{s_{c-t+1}+\dots+s_c} \binom{k_1-c-1}{s_c-1} \cdots \binom{k_t-c-1}{s_{c-t+1}-1} \\
& \quad = \sum_{i_{d+1}=1}^n \sum_{i_{d+2}=1}^n \cdots \sum_{i_{d+t}=1}^n A(n; s_1, \dots, s_{c-t}; i_1, \dots, i_{d+t}) (-1)^{tn} \\
& \quad \quad \times \binom{k_1-2n+d-2+i_{d+1}}{i_{d+1}-1} \cdots \binom{k_t-2n+d-2+i_{d+t}}{i_{d+t}-1}.
\end{aligned}$$

By the Chu-Vandermonde summation, the right-hand side is equal to

$$\begin{aligned}
& \sum_{i_{d+1}=1}^n \sum_{i_{d+2}=1}^n \cdots \sum_{i_{d+t}=1}^n \sum_{s_c=1}^{i_{d+1}} \sum_{s_{c-1}=1}^{i_{d+2}} \cdots \sum_{s_{c-t+1}=1}^{i_{d+t}} A(n; s_1, \dots, s_{c-t}; i_1, \dots, i_{d+t}) (-1)^{tn} \\
& \quad \times \binom{k_1-c-1}{s_c-1} \binom{-2n+d-1+i_{d+1}+c}{i_{d+1}-s_c} \binom{k_2-c-1}{s_{c-1}-1} \binom{-2n+d-1+i_{d+2}+c}{i_{d+2}-s_{c-1}} \\
& \quad \quad \cdots \binom{k_t-c-1}{s_{c-t+1}-1} \binom{-2n+d-1+i_{d+t}+c}{i_{d+t}-s_{c-t+1}}.
\end{aligned}$$

Consequently,

$$A(n; s_1, \dots, s_c; i_1, \dots, i_d) = \sum_{i_{d+1}=s_c}^n \sum_{i_{d+2}=s_{c-1}}^n \dots \sum_{i_{d+t}=s_{c-t+1}}^n A(n; s_1, \dots, s_{c-t}; i_1, \dots, i_{d+t}) \\ \times (-1)^{s_{c-t+1}+\dots+s_c+tn} \binom{-2n+d-1+i_{d+1}+c}{i_{d+1}-s_c} \dots \binom{-2n+d-1+i_{d+t}+c}{i_{d+t}-s_{c-t+1}}.$$

Equivalently, by using  $\binom{n}{k} = (-1)^k \binom{k-n-1}{k}$ ,

$$A(n; s_1, \dots, s_c; i_1, \dots, i_d) = \sum_{i_{d+1}=s_c}^n \sum_{i_{d+2}=s_{c-1}}^n \dots \sum_{i_{d+t}=s_{c-t+1}}^n A(n; s_1, \dots, s_{c-t}; i_1, \dots, i_{d+t}) \\ \times (-1)^{i_{d+1}+\dots+i_{d+t}+tn} \binom{2n-c-d-s_c}{i_{d+1}-s_c} \dots \binom{2n-c-d-s_{c-t+1}}{i_{d+t}-s_{c-t+1}}. \quad (5.1)$$

In the special case  $t = 1$ ,  $c = 2$  and  $d = 0$ , this gives

$$A(n; s_1, s_2; -) = \sum_{i_1=s_2}^n (-1)^{n+i_1} \binom{2n-2-s_2}{i_1-s_2} A(n; s_1; i_1). \quad (5.2)$$

We fix the following notation:  $A_n := A(n; -, -)$ ,  $A_{n,i} := A(n; i; -)$ ,  $B_{n,i,j} = A(n; i; j)$  and  $A_{n,i,j} = A(n; i; j; -)$ . That is,  $A_n$  is the total number of  $n \times n$  alternating sign matrices,  $A_{n,i}$  is the number of  $n \times n$  alternating sign matrices where the unique 1 in the bottom (equivalently top) row is in column  $i$ ,  $B_{n,i,j}$  is the number of  $n \times n$  alternating sign matrices where the unique 1 in the top row is in column  $j$  and the unique 1 in the bottom row is in column  $i$  and, finally,  $A_{n,i,j}$  is the number of monotone triangles with bottom row  $(1, 2, \dots, i-1, i+1, \dots, j-1, j+1, \dots, n)$ . (Or, equivalently,  $A_{n,i,j}$  is the number of monotone  $(2, n)$ -trapezoids with top row  $(i, j)$  and bottom row  $(1, 2, \dots, n)$ . In the introduction, an interpretation of  $A_{n,i,j}$  in terms of alternating sign matrices is provided.)

Stroganov [8, Formula (34)] has shown that

$$B_{n,i,j} - B_{n,i-1,j-1} = \frac{1}{A_{n-1}} (A_{n-1,i-1} (A_{n,j} - A_{n,j-1}) + A_{n-1,j-1} (A_{n,i} - A_{n,i-1})).$$

(See [2] for generalizations of this result.) By the combinatorial interpretation of the numbers  $B_{n,i,j}$  we have  $B_{n,1,j} = A_{n-1,j-1}$  and thus

$$B_{n,i,k} = A_{n-1,k-i} + \sum_{l=2}^i (B_{n,l,k-i+l} - B_{n,l-1,k-i+l-1}) \\ = A_{n-1,k-i} + \frac{1}{A_{n-1}} \sum_{l=2}^i (A_{n-1,l-1} (A_{n,k-i+l} - A_{n,k-i+l-1}) + A_{n-1,k-i+l-1} (A_{n,l} - A_{n,l-1})).$$

If we define  $A_{n,l} = 0$  for  $l \notin \{1, 2, \dots, n\}$  then

$$B_{n,i,k} = \frac{1}{A_{n-1}} \sum_{l=1}^i (A_{n-1,l-1} (A_{n,k-i+l} - A_{n,k-i+l-1}) + A_{n-1,k-i+l-1} (A_{n,l} - A_{n,l-1})).$$

On the other hand, by (5.2), we have

$$A_{n,i,j} = \sum_{k=j}^n (-1)^{n+k} \binom{2n-2-j}{k-j} B_{n,i,k}. \quad (5.3)$$

This finally implies (1.3).

## 6. REMARKS

In this paper we have shown that the number  $A(n; s_1, \dots, s_c; i_1, \dots, i_d)$  of monotone  $(d, n - c)$ -trapezoids with prescribed top and bottom row (or, equivalently, the number of certain  $(n - c - d, n)$ -partial alternating sign matrices) appears as a coefficient of a specialization of  $\alpha(n; k_1, \dots, k_n)$  with respect to a certain (binomial) polynomial basis. We have used this to provide a formula for  $A(n; s_1, s_2; -)$ . Formulas for  $A(n; s_1; -)$  and  $A(n; s_1; i_1)$  were previously known. This raises the question of whether there exist similar formulas for the numbers  $A(n; s_1, \dots, s_c; i_1, \dots, i_d)$  if  $c + d \geq 3$ .

Formula (5.1) shows that it suffices to restrict our attention to the coefficients

$$A(n; -, i_1, \dots, i_d).$$

Next we (re-)derive a system of linear equations for these numbers. (This computation has already appeared in [5].) To this end we need a further identity for  $\alpha(n; k_1, \dots, k_n)$ : observe that there is a one-to-one correspondence between the number of monotone triangles with bottom row  $(k_1, \dots, k_n)$  and the number of monotone triangles with bottom row  $(-k_n, -k_{n-1}, \dots, -k_1)$  and therefore

$$\alpha(n; k_1, \dots, k_n) = \alpha(n; -k_n, -k_{n-1}, \dots, -k_1).$$

Consequently,

$$\begin{aligned} \alpha(n; 1, 2, \dots, n-d, k_{n-d+1}, k_{n-d+2}, \dots, k_n) \\ &= \alpha(n; -k_n, -k_{n-1}, \dots, -k_{n-d+1}, -n+d, -n+d+1, \dots, -1) \\ &= (-1)^{dn-n} \alpha(n; -n+d, -n+d+1, \dots, -1, -k_n-n, -k_{n-1}-n, \dots, -k_{n-d+1}-n) \\ &= (-1)^{dn-n} \alpha(n; 1, 2, \dots, n-d, -k_n-d+1, -k_{n-1}-d+1, \dots, -k_{n-d+1}-d+1). \end{aligned}$$

In terms of the expansion (1.2), this means that

$$\begin{aligned} \sum_{i_1=1}^n \dots \sum_{i_d=1}^n A(n; -, i_1, \dots, i_d) \prod_{l=1}^d \binom{k_{n-d+l} - n + d - 2 + i_l}{i_l - 1} \\ = (-1)^{dn-d} \sum_{j_1=1}^n \dots \sum_{j_d=1}^n A(n; -, j_1, \dots, j_d) \prod_{l=1}^d \binom{-k_{n-d+l} - n - 1 + j_{d+1-l}}{j_{d+1-l} - 1}. \end{aligned}$$

By the Chu–Vandermonde summation, we have

$$\binom{-k_{n-d+l} - n - 1 + j_{d+1-l}}{j_{d+1-l} - 1} = \sum_{i_l=1}^{j_{d+1-l}} \binom{k_{n-d+l} - n + d - 2 + i_l}{i_l - 1} (-1)^{i_l-1} \binom{-2n-1+j_{d+1-l}+d}{j_{d+1-l}-i_l}$$

and, consequently,

$$A(n; -, i_1, \dots, i_d) = \sum_{j_1=i_d}^n \dots \sum_{j_d=i_1}^n (-1)^{dn+i_1+\dots+i_d} A(n; -, j_1, \dots, j_d) \prod_{l=1}^d \binom{-2n-1+j_{d+1-l}+d}{j_{d+1-l}-i_l}.$$

Equivalently, by using  $\binom{n}{k} = (-1)^k \binom{k-n-1}{k}$ ,

$$A(n; -, i_1, \dots, i_d) = \sum_{j_1=i_1}^n \dots \sum_{j_d=i_d}^n (-1)^{dn+j_1+\dots+j_d} A(n; -, j_d, \dots, j_1) \prod_{l=1}^d \binom{2n-i_l-d}{j_l-i_l} \quad (6.1)$$

where  $1 \leq i_1, i_2, \dots, i_d \leq n$ . (This is Conjecture 5 from [5].)

**Remark 1.** In fact, we have used the principal that certain identities for  $\alpha(n; k_1, \dots, k_n)$  translate to identities for the coefficients  $A(n; s_1, \dots, s_c; i_1, \dots, i_d)$ . Another (much less interesting) example in this respect is

$$\begin{aligned} \alpha(n; k_1, \dots, k_c, c+1, \dots, n-d, k_{n-d+1}, \dots, k_n) \\ = \alpha(n; -k_n, -k_{n-1}, \dots, -k_{n-d+1}, -n+d, \dots, -c-1, -k_c, \dots, -k_1) \\ = \alpha(n; n+1-k_n, n+1-k_{n-1}, \dots, n+1-k_{n-d+1}, d+1, \dots, n-c, n+1-k_c, \dots, n+1-k_1), \end{aligned}$$

which translates in a similar way to

$$A(n; s_1, \dots, s_c; i_1, \dots, i_d) = A(n; i_1, \dots, i_d; s_1, \dots, s_c),$$

an identity that is of course (almost) obvious from the combinatorial interpretation if  $s_1 < s_2 < \dots < s_c$  and  $i_1 < i_2 < \dots < i_d$ .

For fixed  $n$ , the system of linear equations (6.1) provides a total of  $n^d$  linear equations for the  $n^d$  numbers  $A(n; -, i_1, \dots, i_d)$ ,  $1 \leq i_1, i_2, \dots, i_d \leq n$ . However, computer experiments show that there is a certain dependency under these linear equations.

For the case  $d = 1$  (i.e. the numbers  $A(n; -, i) = A_{n,i}$  of the refined alternating sign matrix theorem), it was shown in [3] that (6.1) together with  $A_{n,1} = A_{n-1}$  and the symmetry  $A_{n,i} = A_{n,n+1-i}$  for  $1 \leq i \leq n$  (which easily follows from the combinatorial interpretation of the numbers  $A_{n,i}$ ) provides a system of linear equations that determines the numbers  $A_{n,i}$  uniquely. In [5] we have conjectured that this extends to the case  $d = 2$ , i.e. the numbers  $A(n; -, i, j) = A_{n,i,j}$ . In this case, we have to add  $A_{n,i,n} = A_{n-1,i}$ , the near-symmetry

$$A_{n,i,j} = A_{n,n+1-j,n+1-i}$$

for all  $i, j \in \{1, 2, \dots, n\}$  unless  $(i, j) \notin \{(n-1, 1), (n, 2)\}$  and  $A_{n,n-1,1} - A_{n-1} = A_{n,n,2}$ . If  $i < j$  this near-symmetry also easily follows from the combinatorial interpretation of the numbers  $A_{n,i,j}$ , however, for  $i \geq j$ , we took some efforts in [5] to derive it. It would be of interest to see whether this (conjectural) behaviour extends to the case  $d > 2$ . By Cramer's rule this would then at least provide a determinantal expression for the numbers  $A(n; -, i_1, \dots, i_d)$ .

As mentioned above, monotone  $(d, n-c)$ -trapezoids correspond to certain  $(n-c-d, n)$ -partial alternating sign matrices. The latter are generalizations of alternating sign matrices with loosened columnsum restrictions. Clearly, a next natural step would be to consider even more general partial alternating sign matrices, where we loosen the columnsum restrictions and the rowsum restrictions. In particular, the quadruply refined enumeration of alternating sign

matrices where we fix top and bottom row as well as the first and the last column of the alternating sign matrix, would be of major interest.

ILSE FISCHER

INSTITUT FÜR MATHEMATIK, UNIVERSITÄT KLAGENFURT

9020 KLAGENFURT, AND

FAKULTÄT FÜR MATHEMATIK, UNIVERSITÄT WIEN

1090 WIEN, AUSTRIA

`Ilse.Fischer@univie.ac.at`

## REFERENCES

- [1] D. M. Bressoud, *Proofs and Confirmations, The Story of the Alternating Sign Matrix Conjecture*, Cambridge University Press, Cambridge, 1999.
- [2] F. Colomo and A. G. Pranko, On two-point boundary correlations in the six-vertex model with DWBC, *J. Stat. Mech.* (2005) P 05010.
- [3] I. Fischer, A new proof of the refined alternating sign matrix theorem. *J. Comb. Theory Ser. A* 114 (2007), 253–264.
- [4] I. Fischer, The operator formula for monotone triangles – simplified proof and three generalizations, [arXiv:0903.4628](#).
- [5] I. Fischer and D. Romik, More refined enumerations of alternating sign matrices, [arXiv:arXiv:0903.5073](#).
- [6] M. Karklinsky and D. Romik, A formula for the doubly refined enumeration of alternating sign matrices, [arXiv:0906.3405v1](#).
- [7] W.H. Mills, D.P. Robbins, H. Rumsey, Alternating sign matrices and descending plane partitions. *J. Comb. Theory Ser. A* 34 (1983), 340–359.
- [8] Yu. G. Stroganov, A new way to deal with Izergin-Korepin determinant at root of unity, [arXiv:math-ph/0204042](#).
- [9] D. Zeilberger, Proof of the alternating sign matrix conjecture. *Electronic J. Combin* 3 (1996), R13.